


-continued




[1233] 3-cyano-N'-(4-[(cyclohexylmethyl)oxy]phenyl)methyl)-N'-(2-hydroxyethyl)-4-[(1-methyl-ethyl)oxy]benzohydrazide


[1233] Following the procedure of Example 116b), except substituting the compound from Example 118a) for the compound from Example 114a), the title compound was obtained as a glassy solid. ^1H NMR (400 MHz, DMSO- d_6) δ ppm 9.37 (s, 1H) 7.99 (d, J =2.3 Hz, 1H) 7.95 (dd, J =8.8, 2.3 Hz, 1H) 7.33 (d, J =9.1 Hz, 1H) 7.25 (d, J =8.8 Hz, 2H) 6.82 (d, J =8.8 Hz, 2H) 4.86 (qq, J =6.1 Hz, 1H) 4.41 (t, J =6.1 Hz, 1H) 3.98 (s, 2H) 3.71 (d, J =6.6 Hz, 2H) 3.44 (q, J =6.0 Hz, 2H) 2.90 (t, J =5.8 Hz, 2H) 1.77 (d, J =12.4 Hz, 2H) 1.58-1.72 (m, 4H) 1.31 (d, J =6.1 Hz, 6H) 1.12-1.28 (m, 3H) 1.03 (dd, J =12.5, 2.0 Hz, 1H) 0.95-1.00 (m, J =11.1, 2.5 Hz, 1H). MS(ES+) m/e 466 [M+H] $^+$.

Example 120

[1234]



-continued



3-chloro-N'-(4-[(2-(1,1-dimethylethyl)imidazol-4-yl)phenyl]methyl)-N'-(3-hydroxypropyl)-4-[(1-methylethyl)oxy]benzohydrazide

[1235] Following the procedure of Example 116b), except substituting the compound from Example 101b) for the compound from Example 116a) and 3-(t-butyldimethylsilyloxy)propanal (prepared by the method of Berque, I.; Le Ménez, P.; Razon, P.; Mahuteau, J.; Férezou, J.-P.; Pancrazi, A.; Ardisson, J.; Brion, J.-D., *J. Org. Chem.*, 1999, 64, 373-381) for (t-butyldimethylsilyloxy)acetaldehyde, the title compound was obtained as a white solid. ^1H NMR (400 MHz, CHLOROFORM-d) δ ppm 9.24 (s, 1H) 7.76 (d, J =2.3 Hz, 1H) 7.64 (dd, J =8.6, 2.3 Hz, 1H) 7.58 (d, J =8.3 Hz, 2H) 7.41 (s, 1H) 7.30 (d, J =8.3 Hz, 2H) 7.19 (d, J =9.1 Hz, 1H) 4.74 (qq, J =6.1 Hz, 1H) 4.45 (t, J =5.2 Hz, 1H) 3.97 (s, 2H) 3.73 (s, 3H) 3.48 (q, J =6.1 Hz, 2H) 2.91 (t, J =6.8 Hz, 2H) 1.53-1.67 (m, 2H) 1.37 (s, 9H) 1.28 (d, J =6.1 Hz, 6H). MS(ES+) m/e 513 [M+H] $^+$.

Example 121

[1236]

